







A Summary of the Gas Laws

Gas Law	<i>Charles' Law</i>	<i>Boyle's Law</i>																
Variables Involved	Volume, Temperature of a gas in Kelvin	Pressure and volume																
What is Constant?	Number of moles and pressure	Number of moles and temperature																
Formula	$\frac{V_1}{T_1} = \frac{V_2}{T_2}$	$P_1V_1 = P_2V_2$																
Graph																		
Qualitative Representation	At constant pressure, a gas' volume is directly proportional to the absolute (Kelvin) temperature.	At constant temperature, a gas' volume is inversely proportional to its pressure.																
Data Example	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>T(K)</th> <th>V(L)</th> </tr> </thead> <tbody> <tr> <td>0.00</td> <td>0.00</td> </tr> <tr> <td>150</td> <td>22</td> </tr> <tr> <td>300.</td> <td>44</td> </tr> </tbody> </table>	T(K)	V(L)	0.00	0.00	150	22	300.	44	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>V(L)</th> <th>P(kPa)</th> </tr> </thead> <tbody> <tr> <td>10.0</td> <td>100.</td> </tr> <tr> <td>20.0</td> <td>50.0</td> </tr> <tr> <td>40.0</td> <td>25.0</td> </tr> </tbody> </table>	V(L)	P(kPa)	10.0	100.	20.0	50.0	40.0	25.0
T(K)	V(L)																	
0.00	0.00																	
150	22																	
300.	44																	
V(L)	P(kPa)																	
10.0	100.																	
20.0	50.0																	
40.0	25.0																	
Molecular Representation																		

Gas Law	Gay Lussac's Law	Avogadro's Law (only one way of representing it)																
Variables Involved	Pressure, Temperature of a gas in Kelvin	Moles and volume																
What is Constant?	Number of moles and volume	Pressure and Temperature																
Formula	$\frac{P_1}{T_1} = \frac{P_2}{T_2}$	$\frac{V_1}{n_1} = \frac{V_2}{n_2}$																
Graph																		
Qualitative Representation	At constant volume, a gas' pressure is directly proportional to the absolute (Kelvin) temperature.	At constant temperature and pressure, a gas' volume is directly proportional to the number of moles, regardless of the type of ideal gas.																
Data Example	<table border="1" data-bbox="630 1096 993 1251"> <thead> <tr> <th>T(K)</th> <th>P(kPa)</th> </tr> </thead> <tbody> <tr> <td>0.00</td> <td>0.00</td> </tr> <tr> <td>150</td> <td>100</td> </tr> <tr> <td>300.</td> <td>200</td> </tr> </tbody> </table>	T(K)	P(kPa)	0.00	0.00	150	100	300.	200	<table border="1" data-bbox="1019 1096 1383 1251"> <thead> <tr> <th>moles</th> <th>V(L)</th> </tr> </thead> <tbody> <tr> <td>1.00</td> <td>22.4</td> </tr> <tr> <td>2.00</td> <td>44.8</td> </tr> <tr> <td>4.00</td> <td>89.6</td> </tr> </tbody> </table>	moles	V(L)	1.00	22.4	2.00	44.8	4.00	89.6
T(K)	P(kPa)																	
0.00	0.00																	
150	100																	
300.	200																	
moles	V(L)																	
1.00	22.4																	
2.00	44.8																	
4.00	89.6																	
Molecular Representation	